arXiv:math/0502134v1 [math.NT] 7 Feb 2005

q-ANALOGUE OF EULER-BARNES' NUMBERS AND POLYNOMIALS

Taekyun Kim and Leecha Jang

Abstract. Recently Kim [2,6] has introduced an interesting Euler-Barnes' numbers and polynomials. In this paper, we construct the q-analogue of Euler-Barnes'numbers and polynomials, and investigate their some properties.

§1. Introduction

Let $w, a_1, a_2, \cdots, a_r$ be complex numbers such that $a_i(\neq 0)$ for each $i, i = 1, 2, \cdots, r$. Then the Euler-Barnes' polynomials of w with parameters $a_1, a_2, \cdots, a_r$ are defined as

$$\frac{(1-u)^r}{\prod_{j=1}^{r}(e^{a_j t}-u)}\, e^{wt} = \sum_{n=0}^{\infty} H_n^{(r)}(w, u|a_1, a_2, \cdots, a_r)\frac{t^n}{n!},$$

for $u \in \mathbb{C}$ with $|u| > 1$, cf.[6]. In the special case $w = 0$, the above polynomials are called the r-th Euler-Barnes' numbers. We write

$$H_n^{(r)}(u|a_1, a_2, \cdots, a_r) = H_n^{(r)}(0, u|a_1, a_2, \cdots, a_r).$$

Throughout this paper, the symbols $\mathbb{Z}$, $\mathbb{Z}_p$, $\mathbb{Q}_p$, $\mathbb{C}$ and $\mathbb{C}_p$ will respectively denote the ring of rational integers, the ring of p-adic integers, the field of p-adic numbers, the complex number field and the completion of algebraic closure of $\mathbb{Q}_p$. Let ν_p be the normalized exponential valuation of $\mathbb{C}_p$ with $|p|_p = p^{-\nu_p(p)} = p^{-1}$. When one talks of q-extension, q is variousely considered as an indeterminate, a complex number $q \in \mathbb{C}$,

---

1991 *Mathematics Subject Classification.* 11B68.

Key words and phrases. Euler numbers, Bernoulli numbers, zeta function.

Typeset by $\mathcal{AMS}$-TEX

or p-adic number $q \in \mathbb{C}_p$, cf.[2-5]. If $q \in \mathbb{C}$, one normally assumes $|q| < 1$. If $q \in \mathbb{C}_p$, one normally assumes $|1-q|_p \le p^{-\frac{1}{p-1}}$, so that $q^x = exp(x \log q)$. In this paper we use the notation:

$$[x] = [x:q] = \frac{1-q^x}{1-q},\ [x:z] = \frac{1-z^x}{1-z},\ \text{ cf. } [1,2,8].$$

The ordinary Euler numbers E_m are defined by the generating function in the complex number field as

$$\frac{2}{e^t+1} = \sum_{m=0}^{\infty} E_m \frac{t^m}{m!},\ \ (|t| < \pi),\ \text{ cf. } [9].$$

Let u be an algebraic in complex number field. Then Frobenius-Euler numbers are defined as

$$\frac{1-u}{e^t-u} = \sum_{n=0}^{\infty} H_n(u) \frac{t^n}{n!},\ \ (|t| < \pi),\ \text{ cf. } [9,10].$$

Note that $H_n(-1) = E_n$. Also, Carlitz defined the q-analogue of Frobenius-Euler numbers and polynomials as follows:

$$H_0(u:q) = 1, (qH+1)^k - uH_k(u:q) = 0, \text{ if } k \ge 1,$$

where u is a complex number with $|u| > 1$:

$$H_k(u, x:q) = (q^x H + [x])^k, \text{ if } k \ge 0,\ \text{ cf. } [2,11],$$

with the usual convention about replacing $H^k(u:q)$ by $H_k(u:q)$. For any positive integer N, $z \in \mathbb{C}_p$,

$$\mu_z(a + p^N\mathbb{Z}_p) = \frac{z^a}{[p^N:z]}$$

can be extended to distribution on $\mathbb{Z}_p$, cf. [1,2,7,13]. Let $UD(\mathbb{Z}_p)$ be denoted by the set of uniformly differentiable functions on $\mathbb{Z}_p$. Then this distribution admits the following integral for $f \in UD(\mathbb{Z}_p)$:

$$I_z(f) = \int_{\mathbb{Z}_p} f(x) d\mu_z(x) = \lim_{N\to\infty} \frac{1}{[p^N:z]} \sum_{x=0}^{p^N-1} f(x) z^x,\ \text{ cf. } [1,2,12].$$

The purpose of this paper is to construct the q-analogue of Euler-Barnes' numbers and investigate their some properties.

§2. q-Analogue of multiple Euler numbers and polynomials

Let d be a fixed integer and let p be a fixed prime number. We set

$$\begin{aligned} X &= \varprojlim_{N} (\mathbb{Z}/dp^N\mathbb{Z}), \\ X^* &= \bigcup_{\substack{0<a<dp \\ (a,p)=1}} a + dp\mathbb{Z}_p, \\ a + dp^N\mathbb{Z}_p &= \{x \in X \mid x \equiv a \pmod{dp^N}\}, \end{aligned}$$

where $a \in \mathbb{Z}$ lies in $0 \le a < dp^N$.

Let $u \in \mathbb{C}_p$ with $|1-u^f|_p \ge 1$ for each positive integer f and let $a_1, a_2, \cdots, a_r$ be non-zero p-adic integers. For $w \in \mathbb{Z}_p$, we consider the q-analogue of Euler-Barbes' polynomials by using p-adic invariant integrals as follows: For $q \in \mathbb{C}_p$ with $|1-q|_p < p^{-\frac{1}{1-p}}$, define

$$(1) \quad H_n^{(r)}(w,u,q|a_1,a_2,\cdots,a_r) = \underbrace{\int_{\mathbb{Z}_p} \cdots \int_{\mathbb{Z}_p}}_{r \text{ times}} [w + \sum_{j=1}^r a_j x_j : q]^n d\mu_u(x_1) \cdots d\mu_u(x_r).$$

By (1), we note that

$$\begin{aligned} &\underbrace{\int_{\mathbb{Z}_p} \cdots \int_{\mathbb{Z}_p}}_{r \text{ times}} [w + \sum_{j=1}^r a_j x_j : q]^n d\mu_u(x_1) \cdots d\mu_u(x_r) \\ &= \lim_{N\to\infty} \frac{1}{[p^N : u]^r} \sum_{x_1,\cdots,x_r=0}^{p^N-1} [w + \sum_{j=1}^r a_j x_j : q]^n u^{\sum_{j=1}^r x_j} \\ &= \lim_{N\to\infty} \left(\frac{1-u}{1-u^{p^N}}\right)^r \sum_{x_1,\cdots,x_r=0}^{p^N-1} \left(\sum_{l=0}^n \binom{n}{l} (\frac{1}{1-q})^n (-1)^l q^{l(w+\sum_{j=1}^r a_j x_j)} u^{\sum_{j=1}^r x_j} \right) \\ &= \frac{(1-u)^r}{(1-q)^n} \sum_{l=0}^n \binom{n}{l} (-1)^l q^{lw} \left(\frac{1}{\prod_{j=1}^r (1-q^{la_j}u)} \right), \end{aligned}$$

where $\binom{n}{l}$ is binomial coefficient. Therefore we obtain the following:

Theorem 1. *For $n \geq 0$, we have*

$$H_n^{(r)}(w,u,q|a_1,\cdots,a_r) = \frac{(1-u)^r}{(1-q)^n}\sum_{l=0}^{n}\binom{n}{l}(-1)^l q^{lw}\left(\frac{1}{\prod_{j=1}^{r}(1-q^{la_j}u)}\right).$$

Moreover,

$$\lim_{q\to 1} H_n^{(r)}(w,u,q|a_1,\cdots,a_r) = H_n^{(r)}(w,u^{-1}|a_1,\cdots,a_r).$$

Remark. *(1) In the special case $w=0$, we write*

$$H_n^{(r)}(u,q|a_1,\cdots,a_r) = H_n^{(r)}(0,u,q|a_1,\cdots,a_r).$$

(2) Note that $\lim_{q\to 1} H_n^{(1)}(u,q|1) = H_n(u^{-1})$, *cf.[8,9].*

Let $G_q^{(r)}(t,u|a_1,a_2,\cdots,a_r)$ be the generating function of $H_n^{(r)}(u,q|a_1,\cdots,a_r)$:

$$G_q^{(r)}(t,u|a_1,\cdots,a_r) = \sum_{k=0}^{\infty} H_k^{(r)}(u,q|a_1,\cdots,a_r)\frac{t^k}{k!},$$

for $q \in \mathbb{C}_p$ with $|1-q|_p < 1$, $u \in \mathbb{C}_p$ with $|1-u^f|_p \geq 1$. Then we have

$$\begin{aligned}
&G_q^{(r)}(t,u|a_1,\cdots,a_r)\\
&= \sum_{k=0}^{\infty} H_k^{(r)}(u,q|a_1,\cdots,a_r)\frac{t^k}{k!}\\
&= \sum_{k=0}^{\infty}\frac{(1-u)^r}{(1-q)^k}\sum_{i=0}^{k}\binom{k}{i}(-1)^i\left(\prod_{l=1}^{r}\frac{1}{1-q^{ia_l}u}\right)\frac{t^k}{k!}\\
&= (1-u)^r e^{\frac{t}{1-q}}\sum_{j=0}^{\infty}\left(\prod_{l=1}^{r}\frac{1}{1-q^{ja_l}u}\right)\left(\frac{1}{1-q}\right)^j\frac{t^j}{j!}.
\end{aligned}$$

Therefore we obtain the following:

Theorem 2. *For $q \in \mathbb{C}_p$ with $|1-q|_p < 1$, $u \in \mathbb{C}_p$ with $|1-u^f|_p \geq 1$, we have*

$$G_q^{(r)}(t,u|a_1,\cdots,a_r) = e^{\frac{t}{1-q}}(1-u)^r\sum_{j=0}^{\infty}\left(\prod_{l=1}^{r}\frac{1}{1-q^{ja_l}u}\right)\left(\frac{1}{1-q}\right)^j\frac{t^j}{j!}.$$

Corollary 3. *For $q \in \mathbb{C}_p$ with $|1-q|_p < 1$, $u \in \mathbb{C}_p$ with $|1-u|_p \geq 1$, we have*

$$\begin{aligned} & G_q^{(r)}(x,t,u|a_1,\cdots,a_r) \\ &= \sum_{n=0}^{\infty} H_n^{(r)}(x,u,q|a_1,\cdots,a_r)\frac{t^n}{n!} \\ &= e^{\frac{t}{1-q}}(1-u)^r \sum_{j=0}^{\infty}\left(\prod_{l=1}^{r}\frac{1}{1-q^{ja_l}u}\right)\left(\frac{1}{1-q}\right)^j q^{jx}\frac{t^j}{j!}. \end{aligned}$$

Note that

$$\lim_{q\to 1} G_q^{(r)}(x,t,u|a_1,\cdots,a_r) = \frac{(1-u^{-1})^r}{\prod_{l=1}^{r}(e^{a_j t}-u^{-1})}\, e^{xt}.$$

By (1), the Euler-Barnes' polynomials of x can be rewritten as

$$H_n^{(r)}(w,u,q|a_1,\cdots,a_r) = \sum_{k=0}^{n}\binom{n}{k}[w:q]^{n-k}q^{wk}H_k^{(r)}(u,q|a_1,\cdots,a_r).$$

From the above Eq.(1), we have the distribution relation for the q-analogue of Euler-Barnes'polynomials as follows:

Theorem 4. *For $f \in \mathbb{N}$, we have*

$$\begin{aligned} & \frac{1}{(u-1)^r}H_n^{(r)}(w,u,q|a_1,\cdots,a_r) \\ &= [f:q]^n \sum_{i_1,\cdots,i_r=0}^{f-1}\frac{u^{\sum_{j=1}^{r} i_j}}{(u^f-1)^r}H_n^{(r)}(\frac{w+\sum_{j=1}^{r}a_j i_j}{f},u^f,q^f|a_1,\cdots,a_r). \end{aligned} \tag{2}$$

This is equivalent to

$$\begin{aligned} & \frac{1}{(u-1)^r}H_n^{(r)}(w,u,q|a_1,\cdots,a_r) \\ &= [f:q]^n \sum_{i_1,\cdots,i_r=0}^{f-1}\frac{u^{\sum_{j=1}^{r} i_j}}{(u^f-1)^r}H_n^{(r)}(\frac{w+\sum_{j=1}^{r}a_j i_j}{f},u^f,q^f|a_1,\cdots,a_r). \end{aligned}$$

For $k \geq 0$, $f \in \mathbb{N}$, we set

$$E^{(k)}_{u:a_1,q}(x + fp^k\mathbb{Z}_p) = \frac{[fp^N : q]^k u^x}{1 - u^{fp^N}} H^{(1)}_k\left(\frac{a_1 x}{fp^N}, u^{fp^N}, q^{fp^N} | a_1\right), \tag{3}$$

and this can be extended to a distribution on X. We show that $E^{(k)}_{u:a_1,q}$ is a distribution on X. For this, it suffices to check that

$$\sum_{i=0}^{p-1} E^{(k)}_{u:a_1,q}(x + ifp^N + fp^{N+1}\mathbb{Z}_p) = E^{(k)}_{u:a_1,q}(x + fp^k\mathbb{Z}_p).$$

By (2), we easily see that

$$\begin{aligned}
&\sum_{i=0}^{p-1} \frac{[p : q^{fp^N}]^k}{1 - (u^{fp^N})^p} (u^{fp^N})^i H^{(1)}_k\left(\frac{\frac{a_1 x}{fp^N} + ia_1}{p}, (u^{fp^N})^p, (q^{fp^N})^p | a_1\right) \\
&= \frac{1}{1 - u^{fp^N}} H^{(1)}_k\left(\frac{a_1 x}{fp^N}, u^{fp^N}, q^{fp^N} | a_1\right).
\end{aligned}$$

Therefore, we have

$$\begin{aligned}
&\sum_{i=0}^{p-1} E^{(k)}_{u:a_1,q}(x + ifp^N + fp^{N+1}\mathbb{Z}_p) \\
&= \sum_{i=0}^{p-1} \frac{[fp^{N+1} : q]^k u^{(x+ifp^N)}}{1 - u^{fp^{N+1}}} H^{(1)}_k\left(\frac{a_1(x + ifp^N)}{fp^{N+1}}, u^{fp^{N+1}}, q^{fp^{N+1}} | a_1\right) \\
&= u^x \sum_{i=0}^{p-1} \frac{[fp^N : q]^k [p : q^{fp^N}]^k}{1 - (u^{fp^N})^p} (u^{fp^N})^i H^{(1)}_k\left(\frac{\frac{a_1 x}{fp^N} + ia_1}{p}, (u^{fp^N})^p, (q^{fp^N})^p | a_1\right) \\
&= \frac{u^x [fp^N : q]^k}{1 - u^{fp^N}} H^{(1)}_k\left(\frac{a_1 x}{fp^N}, u^{fp^N}, q^{fp^N} | a_1\right) \\
&= E^{(k)}_{u:a_1,q}(x + fp^k\mathbb{Z}_p).
\end{aligned} \tag{4}$$

Next we show that $|E^{(k)}_{u:a_1,q}|_p \leq 1$. Indeed,

$$\begin{aligned}
&E^{(k)}_{u:a_1,q}(x + fp^N\mathbb{Z}_p) \\
&= \sum_{i=0}^{k} \binom{k}{i} \left(\frac{u^x}{1 - u^{fp^N}}\right) [a_1 x : q]^{k-i} [fp^N : q]^i q^{a_1 x i} H^{(1)}_i(u^{fp^N}, q^{fp^N} | a_1).
\end{aligned} \tag{5}$$

By induction on i, we see that

$$\left|\frac{u^x}{1-u^{fp^N}}H_i^{(1)}(u^{fp^N}, q^{fp^N}|a_1)\right|_p \leq 1, \text{ for all } i,$$

where we use the assumption $|1-u^f|_p \geq 1$, it follows that we have

$$|E_{u:a_1,q}^{(k)}(x+ifp^N+fp^N\mathbb{Z}_p)|_p \leq 1. \tag{6}$$

Thus $E_{u:a_1,q}^{(k)}$ is a measure on X. This measure yields an integral for each non-negative integers k as follows:

Proposition 5. *For $k \geq 0$, we have*

$$\int_X dE_{u:a_1,q}^{(k)}(x) = \int_{\mathbb{Z}_p} dE_{u:a_1,q}^{(k)} = \frac{1}{1-u}H_k^{(1)}(u,q|a_1).$$

It is easy to see that

$$H_0(u,q|a_1) = 1.$$

We may now mention the following formula which is easy to prove by (5) and (6):

$$E_{u:a_1,q}^{(k)}(x+fp^N\mathbb{Z}_p) = [a_1x:q]^k\frac{u^k}{1-u^{fp^N}} + [fp^N:q]\times(p-\text{integral}).$$

Hence, we obtain the following :

$$\begin{aligned}\int_X dE_{u:a_1,q}^{(k)}(x) &= \frac{1}{1-u}\int_X [a_1x:q]^k d\mu_u(x)\\ &= \frac{1}{1-u}H_k^{(1)}(u,q|a_1).\end{aligned}$$

From the above definition, we have the following:

Theorem 6. *Let $a_1, a_2, \cdots, a_r$ be p-adic integers. Then we obtain:*

$$\begin{aligned}&\left(\frac{1}{1-u}\right)^r H_{k,\chi}^{(r)}(u,q|a_1,\cdots,a_r)\\ &= \frac{1}{(1-u^d)^r}[d:q]^k\sum_{i_1,\cdots,i_r=0}^{d-1}u^{\sum_{j=1}^r i_j}\left(\prod_{j=1}^r\chi(i_j)\right)H_k^{(r)}(\frac{\sum_{j=1}^r a_ji_j}{d}, u^d, q^d|a_1,\cdots,a_r).\end{aligned} \tag{7}$$

Note that

$$\int_X \chi(x) dE^{(k)}_{u:a_1,q}(x) = \frac{1}{1-u} H^{(1)}_{k,\chi}(u,q|a_1). \tag{8}$$

Let ω be denoted as the Teichmuller character mod$p$ (if $p=2$, mod4). For $x \in X^*$, we set

$$< x : q >= \frac{[x:q]}{w(x)}.$$

Note that $| < x : q > -1|_p < p^{-\frac{1}{p-1}}$, $< x : q >^s$ is defined as $\exp(s \log_p < x : q >)$ for $|s|_p \le 1$. For $s \in \mathbb{Z}_p$, define

$$L_{p,q:a_1}(u|s,\chi) = \int_{X^*} < a_1 x : q >^{-s} \chi(x) d\mu_u(x).$$

Then we have

$$\begin{aligned} &\frac{1}{1-u} L_{p,q:a_1}(u:-k,x) \\ &= \frac{1}{1-u} H^{(1)}_{k,\chi}(u,q|a_1) - \frac{\chi(p)[p:q]^k}{1-u^p} H^{(1)}_{k,\chi}(u^p,q^p|a_1). \end{aligned}$$

Indeed,we see

$$\begin{aligned} &\int_{X^*} < a_1 x : q >^k \chi\omega^k(x) d\mu_u(x) \\ &= \int_X \chi(x)[a_1 x : q]^k d\mu_u(x) - \chi(p)[p:q]^k \frac{1-u}{1-u^p} \int_X [a_1 x : q^p]^k d\mu_{u^p}(x). \end{aligned}$$

Since $| < a_1 x : q > -1|_p < p^{-\frac{1}{p-1}}$ for $x \in X^*$, we obtain

$$< a_1 x : q >^{p^n} \equiv 1 \pmod{p^n}.$$

For $k \equiv k' \pmod{(p-1)p^n}$, we have

$$L_{p,q:a_1}(u:-k,\chi\omega^k) \equiv L_{p,q:a_1}(u:-k',\chi\omega^{k'}) \pmod{p^n}.$$

References

1. T. Kim, *An invariant p-adic Integral associated with Daehee numbers*, Integral Transforms and Special Functions **13** (2002), 65-69.
2. T. Kim, *p-adic q-integral associated with Changhee-Barnes' q-Bernoulli polynomials*, Integral Trans. Special Funct. **15** (2004), 00-00.
3. T. Kim, *Kummer Congruence for the Bernoulli numbers of higher order*, Appl. Math. Comput. **151** (2004), 589-593.
4. T. Kim, *q-Riemann Zeta functions*, Int. J. Math.Math.Sci. **2004 no.12** (2004), 599-605.
5. T. Kim, *Analytic continuation of multiple q-Zeta functions and their values at negative integers*, Russian J. Math. Phys. **11** (2004), 71-76.
6. T. Kim, *On Euler-Barnes multiple zeta functions*, Russian J. Math. Phys **10** (2003), 261-267.
7. T. Kim, *q-Volkenborn integration*, Russian J. Math. Phys **9** (2002), 288-299.
8. T. Kim, *On p-adic q-L-functions and sums of powers*, Discrete Math. **252** (2002), 179-187.
9. T. Kim, *Some formulae for the q-Bernoulli and Euler polynomials of higher order*, J. Math. Anal. Appl. **273** (2002), 236-242.
10. T. Kim, *A note on q-multiple Zeta function*, J. Physics **34** (2001), 643-646.
11. T. Kim, *On p-adic q-Bernoulli numbers*, J. Korean Math. Soc. **37** (2000), 27-30.
12. T. Kim, *A note on Dirichlet L-series*, Proc. Jangjeon Math. Soc. **6** (2004), 161-166.
13. T. Kim, *A note on the q-analogue of multiple zeta function*, Advan. Stud. Contemp. Math. **8** (2004), 111-113.

Institute of Science Education, Kongju National University, Kongju 314-701, Korea, tkim@kongju.ac.kr

Department of Mathematics and Computer Science, KonKuk University, Choongju, Chungbuk 380-701, Korea, leechae.jang@kku.ac.kr